

Comment on: **Red blood cell distribution width as a surrogate biomarker of damage and disease activity in patients with systemic lupus erythematosus**

Sirs,

We have read with great interest the recently published article by Mercader-Salvans *et al.* (1). This study evaluated the role of red blood cell distribution width (RDW) as a marker of both damage and disease activity in systemic lupus erythematosus (SLE). As the authors stated, this is the largest study studying the usefulness of this interesting parameter in SLE. Moreover, the authors described the RDW association with the individual items of both the SLEDAI-2K score and the SLICC Damage Index after adjustment, confirming its high value in the setting of SLE.

Our group has extensively studied the role of RDW in inflammatory conditions such as sepsis and severe COVID-19, in addition to SLE (2-4). In parallel, we tried to unveil the molecular mechanisms underlying the increased anisocytosis in a systemic inflammation environment (5). In this line of work, we showed that RDW is an independent prognostic marker of death in septic patients admitted to the ICU, improving SOFA, LODS, APACHE-II and SAPS-II scores discrimination ability. Similarly, we confirmed that RDW values were higher in deceased patients from severe COVID-19 than in survivors, even after adjustment by haemoglobin levels or tocilizumab exposure. In both studies, RDW displayed a better discrimination ability for mortality than other markers such as C-reactive protein, procalcitonin or ferritin. In SLE, likewise the study from Mercader-Salvans *et al.*,

RDW in our smaller cohort was associated with serological activity and correlated with SLEDAI-2K and SLICC/ACR scores (1). However, our investigation also showed that interleukin-6 (IL-6), key in both COVID-19 and SLE pathogenesis, was the determinant of the higher RDW values seen in these settings (5-7). Therefore, in SLE, RDW presented a linear correlation with IL-6 levels after adjustment by age and haemoglobin values. Accordingly, we believe that our research complements the study by Mercader-Salvans *et al.*

Altogether, it seems that RDW could indeed be a reliable parameter monitoring SLE activity due to IL-6 systemic and pleiotropic effects on haematopoiesis (8). However, further studies are needed in order to validate the findings from our colleagues and to understand RDW activity-monitoring ability in patients under immunosuppressant drugs and other haematological conditions.

V. MORENO-TORRES^{1,2}, MD, PhD
R. CASTEJÓN DÍAZ¹, MD, PhD

¹Department of Internal Medicine, Health Research Institute Puerta de Hierro-Segovia de Arana (IDIPHIM) Hospital Universitario Puerta de Hierro Majadahonda, Madrid;
²UNIR Health Sciences School and Medical Center, Madrid, Spain.

Please address correspondence to:
Victor Moreno-Torres
Unidad de Enfermedades Autoinmunes Sistémicas, Servicio de Medicina Interna, Hospital Universitario Puerta de Hierro Majadahonda, Instituto de Investigación Sanitaria Puerta de Hierro-Segovia de Arana, C/ Joaquín Rodrigo nº 2, 28222 Majadahonda (Madrid), Spain.
E-mail: victor.moreno.torres.1988@gmail.com

Competing interests: none declared.

© Copyright CLINICAL AND EXPERIMENTAL RHEUMATOLOGY 2024

References

- MERCADER-SALVANS J, GARCÍA-GONZÁLEZ M, QUEVEDO-ABELEDÓ JC *et al.*: Red blood cell distribution width as a surrogate biomarker of damage and disease activity in patients with systemic lupus erythematosus. *Clin Exp Rheumatol* 2024; 42: 1773-80.
<https://doi.org/10.55563/clinexprheumatol/f0jnnm>
- MORENO-TORRES V, SÁNCHEZ-CHICA E, CASTEJÓN R *et al.*: Red blood cell distribution width as a marker of hyperinflammation and mortality in COVID-19. *Ann Palliat Med* 2022; 11: 2609-21. <https://doi.org/10.21037/apm-22-119>
- MORENO-TORRES V, ROYUELA A, MÚÑEZ-RUBIO E *et al.*: Red blood cell distribution width as prognostic factor in sepsis: A new use for a classical parameter. *J Crit Care* 2022; 71: 154069. <https://doi.org/10.1016/j.jcrc.2022.154069>
- MORENO-TORRES V, CASTEJÓN R, MELLOR-PI-TA S *et al.*: Usefulness of the hemogram as a measure of clinical and serological activity in systemic lupus erythematosus. *J Transl Autoimmun* 2022; 5: 100157. <https://doi.org/10.1016/j.jtauto.2022.100157>
- MORENO-TORRES V, CASTEJÓN R, MARTÍNEZ-URBISTONDO M *et al.*: Serum cytokines to predict systemic lupus erythematosus clinical and serological activity. *Clin Transl Sci* 2022; 15: 1676-86. <https://doi.org/10.1111/cts.13283>
- KAUL A, GORDON C, CROW MK *et al.*: Systemic lupus erythematosus. *Nat Rev Dis Primers* 2016; 2: 16039. <https://doi.org/10.1038/nrdp.2016.39>
- RUIZ-ANTORÁN B, SANCHO-LÓPEZ A, TORRES F *et al.*: Combination of tocilizumab and steroids to improve mortality in patients with severe COVID-19 infection: a Spanish, multicenter, cohort study. *Infect Dis Ther* 2021; 10: 347-62. <https://doi.org/10.1007/s40121-020-00373-8>
- WEISS G, GOODNOUGH LT: Anemia of chronic disease. *New Engl J Med* 2005; 352: 1011-23. <https://doi.org/10.1056/nejmra041809>