



Exploring Internet gaming disorder: an updated perspective of empirical evidence (from 2016 to 2021)

Gemma Mestre-Bach^{a, **}, Fernando Fernandez-Aranda^{b, c, d, e}, Susana Jiménez-Murcia^{b, c, d, e, *}

^a Universidad Internacional de La Rioja, La Rioja, Spain

^b Department of Psychiatry, Bellvitge University Hospital, Barcelona, Spain

^c Departament de Ciències Clíniques, Facultat de Medicina i Ciències de la Salut, Universitat de Barcelona (UB), Feixa Llarga s/n, 08907, Hospitalet de Llobregat, Spain

^d Psychoneurobiology of Eating and Addictive Behaviors Group, Institut d'Investigació Biomèdica de Bellvitge (IDIBELL), 08907 Barcelona, Spain

^e Ciber Fisiopatología Obesidad y Nutrición (CIBEROBN), Instituto de Salud Carlos III, Madrid, Spain

ARTICLE INFO

Keywords:

Online gaming
Addictive behavior
Treatment
Diagnosis
Etiology

ABSTRACT

Background and aims: Since the acceptance of Internet gaming disorder (IGD) as a “disorder due to addictive behaviors”, research has proliferated exponentially. The present review focuses on the conceptualization of IGD, its diagnosis and assessment, associated factors and existing prevention and treatment plans to address it.

Results and conclusions: The discrepancies between the diagnostic criteria for IGD proposed by the two central diagnostic entities, as well as the questioning of their clinical validity, have generated multiple proposals for the diagnosis and psychometric evaluation of IGD. Likewise, there have been numerous suggestions to prevent this pathology, with the involvement of governments, the gaming industry and health institutions. Finally, multiple treatment plans have been proposed, both pharmacological and psychological, although only the efficacy of cognitive behavioral therapy has been tested. It is essential, therefore, to delve deeper into this disorder by addressing the central limitations of the current literature.

1. Introduction

Internet gaming disorder (IGD) was suggested as a mental disorder in section III of the DSM-5 [1] and has been recognized as a “disorder due to addictive behaviors” by the ICD-11 [2], considering it to be a persistent and recurrent pattern of Internet gaming involvement despite the physical and psychological impairment.

However, the conceptualization of this disorder has been a matter for debate. On the one hand, some authors have considered that its inclusion in the diagnostic manuals resulted in the pathologization of some behaviors (in this case, the use of video games), which generated policies and interventions that are not essential; the proposed diagnostic criteria for IGD do not adequately distinguish between highly engaged and problematic gaming behavior, and its addictive nature has also been questioned [3–6]. On the other hand, other authors have emphasized that the diagnosis of IGD is essential, just as it is clear that others, such as anxiety or eating disorders, are essential without the fear of stigmatizing behaviors (which in that case would be worries or eating behaviors) [7].

A recent Delphi study analyzed the criteria proposed by the DSM-5 and ICD-11 to explore a possible consensus among experts about their clinical relevance and diagnostic power [8]. Some criteria (loss of control, gaming despite harm, conflict/inference due to gaming, and functional impairment) were agreed upon because of their clinical usefulness. However, there was also consensus that some DSM-5 criteria (tolerance, deception of others about gaming, and escape/regulating mood) had limited clinical relevance, suggesting its exclusion for the conceptualization and diagnosis of IGD [8].

The most affected by IGD are males (especially adolescents) presenting higher screen hours, craving and negative consequences [9,10]. More specifically, a systematic review that analyzed 53 studies with a total sample of 226,247 individuals highlighted a worldwide IGD prevalence of 3.05% [11]. In the specific case of adolescents, a prevalence of 4.6% has been described [12]. Darvesh et al. [13] found in their review that the prevalence of IGD ranged, in the general population, from 0.21 to 57.50% while in clinical populations it ranged from 3.20 to 91.00%. General population was defined as populations that were not

* Corresponding authors at: University of Barcelona Faculty of Medicine: Universitat de Barcelona Facultat de Medicina i Ciències de la Salut Spain

** Corresponding authors at: Universidad Internacional de La Rioja, La Rioja, Spain

E-mail addresses: gemma.mestre@unir.net (G. Mestre-Bach), ffernandez@bellvitgehospital.cat (F. Fernandez-Aranda), sjimenez@bellvitgehospital.cat, susanajimenez@ub.edu (S. Jiménez-Murcia).

<https://doi.org/10.1016/j.comppsy.2022.152319>

Received 15 October 2021; Received in revised form 7 April 2022; Accepted 13 April 2022

Available online 28 April 2022

0010-440X/© 2022 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

seeking treatment for IGD, did not have IGD at recruitment and were not undergoing treatment for IGD while clinical populations were understood as the ones who sought treatment and those who suffered from IGD at recruitment.

Similarly, Pan et al. [14] conducted a systematic review and meta-analysis of epidemiology of generalized internet addiction and IGD. The authors included 113 epidemiologic studies involving a total of 693,306 individuals, finding a prevalence of 2.47% in the case of IGD.

In view of the lack of scientific consensus regarding IGD, the present narrative review aims to shed light on the assessment, associated factors, prevention, and treatment of this diagnostic category by reviewing studies from the last 5 years. This type of review was selected in order to obtain a broad, critical, and comprehensive perspective of the current knowledge about IGD. For the selection of the studies included in this narrative review, PubMed/Medline was used to search for the term "gaming disorder" and the temporal filter of the last 5 years (September 2016–September 2021). This period was selected due to the proliferation of scientific literature in the field, mainly because of both the acceptance of the condition by ICD-11 and the pandemic period as a consequence of COVID-19.

2. IGD assessment

As highlighted by van Rooij et al. [15], the DSM-5 proposal generated an increase in the design of new psychometric scales to assess IGD. The authors identified six instruments: the 20-item Internet Gaming Disorder Test (IGD-20), a separate nine-item Internet Gaming Disorder Scale–ShortForm (IGDS9-SF), the Internet Gaming Disorder Test (IGDT-10), the Internet Gaming Disorder Scale (IGDS-9), the Clinical-Video Game Addiction Test (C-VAT 2.0) and an unconventional hybrid scale with items pertaining to Internet and online gaming. Similarly, King et al. [16] identified 29 instruments.

In parallel, Bernaldo-de-Quirós et al. [17] found 7 psychometric scales in their systematic review exploring instruments to measure IGD in adolescents and young adults: the IGD-20, the IGDS9-SF, the Problematic Online Gaming Questionnaire (POGQ), the Problematic Online Gaming Questionnaire – Short Form (POGQ-SF), the Video Game Addiction Test (VAT), the C-VAT 2.0 and the IGD scale.

Similarly, King et al. [18], in their comprehensive systematic review, identified 32 instruments to assess IGD, which were included in 320 studies, and highlighted the proliferation of psychometric tools since 2013.

New instruments for the assessment of IGD have been developed, and validations/adaptations of existing instruments have been conducted in the last 5 years (2016–2021) (included in Table 1).

Regarding the geographical origin of the new validated instruments, Asia seems to be the continent most focused on advancing the field of IGD assessment. Moreover, including both new validations and adaptations, China, Korea and Taiwan are the countries with the largest number of studies in this regard. Taking the aims of the studies into account, it seems that IGDS9-SF, IGD-20 and IGDT-10 are the instruments that arouse the greatest scientific interest, given that numerous articles have attempted to test their psychometric properties in different countries. Finally, most of these instruments are based on the criteria proposed by the DSM-5 so they share the same theoretical framework, although the number and type of items vary considerably among them. The shortest tool contains 4 items [68], while the longest is composed of 38 items [48], with the range between 7 and 11 items being the most common. Some authors have reported the need to test the rigor of the IGD diagnostic criteria in multiple populations to increase its accuracy. However, they have pointed out that the existence of multiple IGD assessment instruments greatly hinders scientific progress and that only if the scientific community reaches a consensus and uses exclusively one tool and the same cut-off points can the IGD be fully understood [72].

3. Associated factors

Several theoretical models have attempted to compile all the factors that may be associated with IGD. Among them, the model for Internet addiction [73], the theoretical model of Internet-gaming disorder [74] and the Interaction of Person-Affect-Cognition-Execution (I-PACE) model of specific Internet-use disorders [75] have had special relevance. The different models highlight the relevance of psychopathological factors, personality traits, personal attitudes, cognitive processes, decision-making style, and executive control, among others. The I-PACE model, which encompasses factors proposed by the other two frameworks, proposes: (I) predisposing variables (biopsychological constitution, psychopathological features, personality, and social cognitions, and using motives), (II) affective and cognitive responses to external or internal stimuli (coping, internet-related cognitive biases, cue-reactivity and craving, urge for mood regulation, and attentional biases) and (III) executive functions, inhibitory control and the decision to use certain applications/sites. Thus, multiple factors appear to be associated with the etiology and maintenance of IGD, although further empirical evidence is needed. Some of the factors associated with IGD are summarized below, taking into account the three main blocks considered from the biopsychosocial perspective: biological, psychological and social factors.

3.1. Neurobiological factors

Problematic gaming and IGD have been associated with neural alterations, both in functional and structural studies. More specifically, alterations have been observed in temporoparietal regions (in charge of attentional and self-concept processes), frontal areas (related to cognitive control) and frontal-limbic and subcortical regions (associated with reward processing and emotion regulation processes) [155]. In this vein, notable differences have been observed between individuals with IGD and controls, with the former showing alterations in the neuronal reward system, as well as poor response-inhibition, difficulties in decision-making processes, emotion regulation and cognitive control, poorer working memory and decreased visual and auditory functioning [156–158]. In addition, structural changes such as lower volumes of the striatum and a reduction of the gray-matter volume, as well as the white-matter density, have been described [159,160]. The systematic review by Yao et al. [161] suggests that individuals with IGD present, in comparison with controls: (a) an hypoactivation in the anterior inferior frontal gyrus (in relation to hot-executive function), the posterior insula, somatomotor and somatosensory cortices (in relation to reward function); and (b) an hyperactivation in the caudate, anterior and posterior cingulate cortices, posterior inferior frontal gyrus, (in relation to cold-executive functions and reward).

However, it is worth noting the heterogeneity of the control groups used in these neurobiological studies, given that some use non-gamers, others non-excessive gamers, others amateur gamers, others specify that daily gaming duration was <2 h, and others do not specify the criteria for being a healthy control.

3.2. Psychological factors

3.2.1. Personality and impulsivity

In the last 5 years, the literature on the association between IGD and personality traits, as well as IGD and impulsivity [76–86] has also proliferated. Regarding personality traits, individuals with IGD show maladaptive personality traits, including high levels of negative affectivity and disinhibition [87]. Similarly, an association between IGD and personality traits such as neuroticism, conscientiousness and novelty seeking has also been described [88,89]. Granero et al. [90] have also highlighted the specific role of self-transcendence and cooperation in individuals with IGD, which would entail, among other aspects, a tendency to become immersed in the game context, losing track of time.

Table 1

Main studies in the last 5 years validating/adapting psychometric instruments and clinical interviews.

Authors	Country	Aim	Instrument	Instrument information	Sample
Yu et al. [19]	China	To examine the psychometric properties of a Chinese version of the IGD-20 Test	IGD-20 Test	20-item scale based on IGD items in DSM-5	569 Chinese middle school students and 523 university students
Kim [20]	Korea	To validate the Korean version of the IGD-20 Test.	IGD-20 Test	20-item scale based on IGD items in DSM-5	1,403 Korean online game users
Hawi et al. [21]	Arabia	To validate an Arabic version of the IGD-20 Test.	IGD-20 Test	20-item scale based on IGD items in DSM-5	375 students
Plessis et al. [22]	France	To validate the IGD-20 Test in French.	IGD-20 Test	20-item scale based on IGD items in DSM-5	166 French-speaking video game players who played an average of 21.9 h per week recruited from specialized forums.
Männikkö et al. [23]	Finland	To investigate the psychometric properties of the Finnish version of the IGDT-10	IGDT-10	10-item variant of the IGDT scale	773 Finnish vocational school students
Király et al. [24]	Hungarian, Iranian, English-speaking, French-speaking, Norwegian, Czech, and Peruvian	To test the psychometric properties, including language and gender invariance, in a large international sample of online gamers	IGDT-10	10-item scale assessing IGD as proposed in the DSM-5	7,193 participants comprising international online gamers
Király et al. [25]	Hungary	To develop and validate a brief psychometric instrument (IGDT10)	IGDT-10	10-item scale assessing IGD as proposed in the DSM-5	4,887 gamers (age range 14–64 years old)
Chiu et al. [26]	China	To validate the Chinese version of the IGDT-10	IGDT-10	10-item scale assessing IGD as proposed in the DSM-5	8,110 students from grade four to senior high who played Internet games
Wu et al. [27]	Iran	To verify the construct validity of the Persian IGDS9-SF	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	2,363 Iranian adolescents
Chen et al. [28]	China	To validate the psychometric properties of three simplified Chinese online-related addictive behavior instruments among mainland Chinese primary school students.	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	1108 fourth to sixth graders
Qin et al. [29]	China	To examine the optimal cutoff score for the IGDS9-SF in a Chinese context.	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	Samples from clinical settings (n = 131) and from universities (n = 3,742).
Schivinski et al. [30]	Poland	To develop the Polish version of the IGDS9-SF and scrutinize the nine IGD criteria empirically.	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	3,377 gamers (mean age 20 years old)
Monacis et al. [31]	Italy	To examine the psychometric properties of the IGDS9-SF in an Italian-speaking sample	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	757 participants from Italian schools, universities and gaming halls.
Beranuy et al. [32]	Spain	To translate and adapt the IGDS9-SF to Spanish, as well as to obtain indicators relating to its validity and reliability.	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	535 Vocational Training students who reported playing video games in the past 12 months
Maldonado-Murciano et al. [33]	Spain	To evaluate the psychometric properties of the Spanish version of the IGDS9-SF.	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	388 online gamers
Sánchez-Iglesias et al. [34]	Spain	To assess the psychometric properties of the Spanish version of the IGDS9-SF and criterion validity by analyzing the relationships of their scores with other theoretically related variables.	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	2,173 video game players between 12 and 22 years old
Severo et al. [35]	Brazil	To develop and examine the psychometric properties of the IGDS9-SF in a sample of Brazilian gamers and to find the best cut-off point	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	610 teenagers and adults
Donadon et al. [36]	Brazil	To describe the process of cross-cultural adaptation of the IGDS9-SF to the Brazilian context.	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	30 gamers
Ling et al. [37]	Malaysia	To examine the psychometric properties (factor structure, reliability, and validity) of the Malay Version of the IGDS9-SF.	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	137 Malaysian undergraduates
Evren et al. [38]	Turkey	To test the factor structure, reliability and validity of the IGDS9-SF.	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	1,250 participants
Kim et al. [39]	Korea	To examine the psychometric properties of K-IGDS9-SF.	K-IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	594 Korean game users
Lei et al. [40]	China	To test the psychometric properties of two versions of Chinese IGDSs: a dichotomous IGDS with yes/no type of response and a polytomous IGDS with a 0–5 Likert-type response	short IGDSs	The short IGDSs (9 items) has been developed in two forms with identical items but different response options: a dichotomous IGDS with yes/no type of response; and a polytomous IGDS with a 0–5 Likert-type response	Two population-based samples of Chinese adolescents and adults (351 for dichotomous IGDS and 378 for polytomous IGDS)

(continued on next page)

Table 1 (continued)

Authors	Country	Aim	Instrument	Instrument information	Sample
Sigerson et al. [41]	China	To develop a reliable and valid scale for measuring IGD in Chinese populations	C-IGDS	9-item scale assessing severity of IGD based on IGD items in DSM-5	502 Hong Kong community adults
Paschke et al. [42]	Germany	To investigate the psychometric properties of the IGDS in children and adolescents considering different age and sex groups	Short IGDS	The short IGDSs (9 items)	12- to 17-year old participants. Three independent samples: N = 762, N = 777, and N = 784
Chang et al.	Taiwan	To adjust the Taiwanese version of IGD-SF-T-L based on the DSM-5 criteria to a Likert scale model and test its psychometric property among children and adolescents with ADHD.	IGDS-SF-T-L	A modification of the Taiwanese IGD-SF criteria from a "yes" or "no" model to a 4-point Likert scale	102 children and adolescents with ADHD
van Rooij et al. [43]	Dutch	To develop and test a clinical assessment tool (C-VAT 2.0) in a clinical setting.	C-VAT 2.0	11-item test that covers the proposed DSM-5 criteria	32 clinical young participants (13–23 years old) in treatment for IGD
Lin et al. [44]	Iran	To test the psychometric properties of the Persian GAS through both classical test and modern test theories.	GAS	7-item scale	4,442 adolescents
Liu et al. [45]	China	To examine the psychometric properties of the GAS and to measure invariance among college students in China.	GAS	7-item scale	1040 full-time students from multiple colleges in China
Khazaal et al. [46]	Switzerland	To analyze the GAS in the context of item response theory (graded-response) modeling; and to investigate differential item functioning in 2 subsamples	GAS	7-item scale	2 large representative samples of Swiss men (3320 French-speaking and 2670 German-speaking)
Chen et al. [47]	China	To construct a screening tool to assess IGD symptomatology, defined by the fifth edition of the DSM-5, among Chinese gamers.	C-IGDC	27-item checklist, which is the first multidimensional, DSM-5-based IGD screening tool developed in a Chinese adult population	464 Chinese past-year gamers
Lau et al. [48]	China	To examine the psychometric properties of the IPQ-R for IGD in a general population	IPQ-R for IGD	The 38-item IPQ-R was modified to assess illness representation of IGD	1,501 Chinese community-dwelling adults
Higuchi et al. [49]	Japan	To develop a short screening test for IGD	GAMES test	9-item screening test	281 participants from the general population, aged between 10 and 29 years old.
Yang et al. [50]	China	To develop and validate a checklist for IGD symptoms of Chinese adolescents	DISCA	9 items	2144 secondary school students who reported that they had played Internet games in the past 12 months
Ko et al. [51]	Taiwan	To evaluate the best screening, diagnostic, and prevalence-estimated cutoff points of the CIAS-G for differentiating adults with IGD from regular gamers (without IGD)	CIAS-G	26-item self-report scale to assess participants' online gaming experiences	69 participants with IGD, 69 regular gamers, and 69 healthy participants
Basu et al. [52]	India	To assess IGD using a self-designed scale in conjunction with a screening questionnaire	IGAS	14 items	424 medical students
Irmak et al. [53]	Turkey	To develop a measurement tool based on reporting by family members	GDS-F	18 items that reflect nine diagnostic criteria indicated in Section III of the DSM-5.	173 adolescents from 5th grade to 12th grade, were included in the study
Kim et al. [54]	Korea	To validate the K-MOGQ and to examine the incremental validity of the K-MOGQ beyond the personality variables implicated in IGD	K-MOGQ	27 items that measure motives for online gaming	641 Korean online game users
Wu et al. [55]	China	To empirically evaluate the psychometric properties of the C-MOGQ	C-MOGQ	27 items that measure motives for online gaming	383 Chinese adult online gamers
Yu et al. [56]	China	To investigate the psychometric properties of the Chinese version of the IGCS among adolescents	C-RIGCS	24 items	755 students from junior middle schools
Hawi et al. [57]	Lebanon	To develop and validate an instrument to assess children's overall addiction to digital devices (including video gaming, social media and texting).	DASC	25- item self-report instrument that was developed based on nine diagnostic DSM-5 IGD criteria and also mapped onto Griffiths six core addiction criteria	822 participants from 4th grade to 7th grade
Sarda et al. [58]	France	To assess the reliability and factor structure of the IGD scale, and to provide the cross-validity and criterion validity of this measure.	IGD scale	9 items proposed by Petry et al. referring to their behavior during the last year	693 players
Koo et al. [59]	Korea	To develop and validate the SCI-IGD in adolescents	SCI-IGD	12 items	236 adolescents from both community and clinical settings
Sprong et al. [60]	USA	To compare the DSM-5 validated assessment criteria of the IGD-20 with the primary reinforcing behavioral functions evaluated by the VGFA-R.	IGD-20 and VGFA-R	IGD-20: 20-item scale based on IGD items in DSM-5	304 participants

(continued on next page)

Table 1 (continued)

Authors	Country	Aim	Instrument	Instrument information	Sample
Stavropoulos et al. [61]	Australia, The USA, and The UK	To examine the IGDS9-SF measurement invariance across gamers from Australia, the USA, and the UK	IGDS9-SF	VGFA-R: 24-item Likert-style scale designed to assess four functions that maintain video game play 9-item scale assessing severity of IGD based on IGD items in DSM-5	171 Australian gamers, 463 American gamers, and 281 British gamers
Pontes et al. [62]	The USA, India, and The UK	To examine the IGDS9-SF measurement invariance across gamers from the USA, India, and the UK	IGDS9-SF	9-item scale assessing severity of IGD based on IGD items in DSM-5	1013 gamers from the USA (n = 405), India (n = 336), and the UK (n = 272)
Yilmaz et al. [63]	Turkey	To develop a valid and reliable Video Game Addiction Scale for Children	VASC	21 items	780 children
Jo et al. [64]	Korea	To develop an IGD screening tool for adolescents to be used in various primary care settings.	IGUESS	9 items	121 adolescents recruited from a public school, community-based resources for mental health, a private counseling center, and a psychiatry clinic
Ryu et al. [65]	Korea	To examine the psychometric properties of the DIA in Korean adolescents	DIA	10 items based on the DSM-5 IGD diagnostic criteria	103 adolescents
Besser et al. [66]	Germany	To develop an optimized short screening for problematic Internet use and Internet addiction	CIUS	5 items	3,040 participants
Gmel et al. [67]	Switzerland	To develop a short form of the CIUS	CIUS short form	8 items	learning (n = 1,371) and validation samples (n = 1,550)
Schluter et al. [68]	Canada	To develop a brief self-attribution Screener for Substance and Behavioral Addictions	SSBA	4 items	6,000 adults
Leung et al. [69]	Hong Kong and Taiwan	To investigate whether the BSMAS, SABAS and IGDS-SF9 were appropriate for their use in heterogeneous subsamples from Hong Kong and Taiwan	BSMAS, SABAS, and IGDS-SF9	BSMAS: 6 items to assess social media addiction. SABAS: 6 items to assess smartphone use addiction. IGDS-SF9: 9-item scale assessing severity of IGD based on IGD items in DSM-5	University students from Hong Kong (n = 306) and Taiwan (n = 336)
Chen et al. [70]	Hong Kong and Taiwan	To examine the time invariance of the SABAS, BSMAS, and IGDS-SF9 for Chinese university students, and to evaluate the reproducibility, including relative and absolute reliabilities, of the three instruments across three months	BSMAS, SABAS, and IGDS-SF9	BSMAS: 6 items to assess social media addiction. SABAS: 6 items to assess smartphone use addiction. IGDS-SF9: 9-item scale assessing severity of IGD based on IGD items in DSM-5	640 university students (304 from Hong Kong and 336 from Taiwan)
Wang et al. [71]	34 countries	To evaluate and compare the psychometric properties of the GDT and IGDS9-SF	GDT and IGDS9-SF	GDT: 4-item scale based on the ICD-11 framework. IGDS9-SF: 9-item scale assessing severity of IGD based on IGD items in DSM-5	544 adult gamers

IGD: Internet gaming disorder; IGD-20 Test: Internet Gaming Disorder Test; DSM-5: Diagnostic and Statistical Manual of Mental Disorders- Fifth Edition; IGDT-10: Internet Gaming Disorder Test; IGDS9-SF: Internet Gaming Disorder Scale – Short Form; K-IGDS9-SF: Korean version of the Internet Gaming Disorder Scale—Short Form; IGDSs: Internet Gaming Disorder scales; C-IGDS: Chinese Internet Gaming Disorder Scale; IGDS-SF-T-L: Internet Gaming Disorder Scale with Likert Scale; ADHD: Attention-deficit/hyperactivity disorder; C-VAT 2.0: Clinical Video game Addiction Test; GAS: 7-item Gaming Addiction Scale; C-IGDC: Chinese Internet Gaming Disorder Checklist; C-IGDC: Chinese Internet Gaming Disorder Checklist; IPQ-R: Revised Illness Perception Questionnaire; GAMES test: GAMING Engagement Screener test; DISCA: DSM-5 IGD Symptoms Checklist for Adolescents; CIAS-G: Chen Internet Addiction Scale-Gaming Version; IGAS: Internet gaming addiction scale; GDS-F: Gaming Disorder Scale-Family Form; K-MOGQ: Korean version of the Motives for Online Gaming Questionnaire; C-MOGQ: Chinese version of the Motives for Online Gaming Questionnaire; C-RIGCS: Chinese version of Revised Internet gaming cognition scale; DASC: Digital Addiction Scale for Children; SCI-IGD: Structured Clinical Interview for Internet Gaming Disorder; VGFA-R: Video Game Functional Assessment-Revised; BSMAS: Bergen Social Media Addiction Scale; SABAS: Smartphone Application-Based Addiction Scale; GDT: Gaming Disorder Test; ICD-11: International Classification of Disease, Eleventh Revision; VASC: Videogame Addiction Scale for Children; IGUESS: Internet Game Use-Elicited Symptom Screen; DIA: Diagnostic Interview for Internet Addiction; CIUS: Compulsive Internet Use Scale; SSBA: Screener for Substance and Behavioural Addictions.

However, differences have been observed in individuals with IGD in terms of personality traits and dysfunctionality, establishing two clusters. The first one is characterized by greater introversion, inhibition, borderline tendency traits and lower histrionic traits in comparison to the second cluster which is characterized by less comorbidity and higher functionality [91]. In addition, when comparing individuals with IGD to those with gambling disorder, the former appear to exhibit lower novelty seeking and persistence traits [92].

Regarding impulsivity, a recent systematic review on probability discounting found that individuals with IGD had a greater tendency, compared to controls, to overestimate the potential gains associated

with the consequences of their decisions, especially if these potential gains were not substantial [93]. They also showed a higher tendency to take risks in probabilistic gains. It has also been suggested that subjects with IGD present deficits in decision-making and inhibitory control, aspects that could be associated with brain alterations [94–98]. In this vein, Kräplin et al. [99] highlighted that this impulsive decision-making could be a consequence of IGD, rather than a predictor, due to altered reward learning.

3.2.2. Emotion regulation and emotional intelligence

Another aspect that has been studied in recent years is the role of

emotions in IGD. Specifically, both emotion regulation [100–103] and emotional intelligence [104–106] have been explored. On the one hand, an association has been observed between IGD and impaired emotion regulation, which could exacerbate the negative mood symptoms of young adults with IGD [100,101]. On the other hand, emotional intelligence seems to have a protective effect on the development of IGD [105], and its different dimensions may have different roles in this disorder [104].

3.2.3. Co-occurring symptomatology/disorders

In the last 5 years, the association between IGD and other co-occurring psychopathologies has begun to be studied in depth [107–112], especially since it may increase the severity of the disorder and worsen the treatment outcome. More specifically, the association between IGD and ADHD [113–121], depression [106,122–130], suicidal ideation [131], substance use [132–134], problematic social media use [135,136], sleep problems [135,137,138], autistic traits [103,121,139], anxiety [106,130,140–142], stress [143,144], and posttraumatic stress symptoms [145] has been explored.

The high levels of comorbidity observed in the clinical populations with IGD have been questioned by experts who consider that there is not enough empirical evidence to consider IGD as a mental disorder. They claim that the high levels of comorbidity could reflect that gaming behavior is used as a coping strategy associated with underlying problems that have a different nature [146,147].

3.3. Psychosocial factors

The role of the family in the development and maintenance of problem gaming and IGD seems to be essential [148] and it has been studied from different perspectives: (i) parent status; (ii) the parent-child relationship; (iii) parental influence on gaming behavior; and (iv) family environment [149].

Regarding parental status, factors such as socioeconomic status and parental mental health have been studied, highlighting, for example, an association between IGD and parental anxiety [150]. The parent-child relationship has been the most studied factor, and it has been suggested that those individuals with IGD present greater family conflicts and a poorer family relationship [151], which have even been associated with the severity of the disorder [152]. In addition, over-protection and rejection by parents could also have a negative influence on IGD [153]. Regarding parental influence on gaming, it has been observed that IGD symptomatology may elicit ineffective Internet-specific parental responses that could exacerbate the disorder [154].

4. Policy and prevention

Before considering the proposed policy and prevention plans for IGD, it is worth considering the main harms associated with IGD. Although, compared to other addictive disorders, such as gambling disorder, IGD has relatively low economic costs associated. There is another potential harm. Some common examples associated with extreme gaming time are disturbances in basic activities (such as sleep patterns and fatigue, eating patterns and hygiene), social disturbances and isolation, and neglect of daily responsibilities, as well as physical pain due to poor posture [162].

The main IGD policies applied by the majority of Asian countries have been categorized, in a recent systematic review, into three main blocks: (a) those that limit the availability of games; (b) those that reduce risk and harm, and (c) those that aim to support gamers [163].

In order to limit the availability of games, different policies have been implemented: (i) the shutdown system, in which governments select a time slot and force game service providers to block access to online games during that period of time, for example between 10 p.m. and 6 a.m., as in the case of Thailand; (ii) the fatigue system, in which online game providers have to monitor the playtime of their consumers and discourage minors from playing for long periods of time by reducing

the experience points (XPs) needed to advance in the game associated with the playing time; (iii) parental controls that include content filters, time limits and monitoring the online activity; (iv) limiting gaming time in Internet cafés, for example between 2 p.m. and 8 p.m., as in Thailand; and (v) increasing the price of video games, although this is only a hypothetical measure suggested by the authors, which has not been implemented yet [163].

To reduce risk and harm, some gaming companies are currently implementing warning messages. However, the authors suggest that other potential measures would be to use rating systems for rating games taking into account their addictive potential, as well as to design games that are less addictive or less exploitative of gamers' time [163].

Finally, there are policies to support gamers who develop IGD, although it is not entirely clear which institutions should provide such assistance, whether they should be gaming companies, the government or private institutions [163].

King et al. [164] establish another classification of prevention strategies: (a) universal prevention; (b) selective prevention; and (c) indicated interventions.

Universal prevention may encompass: (i) educational resources that provide gamers with information on healthy levels of use, recommendations on physical activity and promotion of more adaptive activities; (ii) legislative action, such as shutdowns of online gaming services during certain time slots of the day; (iii) technological measures, such as parental controls or notifications for excessive gaming time; (iv) public awareness, such as the stipulation of national days and campaigns that encourage non-use of the Internet; and (v) behavioral measures, such as using the devices in common areas of the home or reducing the number of owned devices [164].

Selective prevention focuses on those subjects who are more vulnerable to IGD, and would consist of: (i) regular screening, for example in schools and universities, in order to identify emerging cases; (ii) mental health checks with medical practitioners to detect possible risk factors; (iii) school-based educational programs, in which healthy patterns of Internet use and gaming are taught; and (iv) workplace Internet policies, with regulations on Internet access [164].

Finally, indicated interventions would be intended for those individuals with IGD, including: (i) support and community groups; (ii) outpatient medical and mental health services; (iii) psychosocial rehabilitation, focused on socialization; and (iv) psychoeducation on strategies to minimize the harm of IGD [164].

Gentile [165] also highlights four central ways in which parents can monitor their children's gaming behavior and, consequently, prevent the development of IGD: (i) co-viewing or co-playing with children or adolescents; (ii) setting restrictions on the amount of time spent on gaming behavior; (iii) setting restrictions on content; and (iv) conducting active mediation with children where perspectives on, for example, the effects of video games are discussed.

As highlighted by Petry et al. [166], at a global level, industries that benefit from products that bring with them harmful consequences, in this case the gaming industry, present a clear conflict of interest when promoting primary, secondary and tertiary prevention strategies, making government regulations and taxation strategies essential. Moreover, the efficacy of all these regulations, as well as their possible side effects, should be systematically tested [167]. However, as Kuss [168] points out, it should be kept in mind that not all prevention policies and plans that are effective in a specific region or cultural context work in other cultures, since the degree of acceptability of these strategies is subject to cross-cultural differences.

4.1. Prevention during COVID-19 pandemic

Video games may have had a positive effect during the lock-downs resulting from the COVID-19 pandemic, as they helped to promote social distancing, socialization and stress reduction, among other factors [169]. However, for the most vulnerable individuals, this increased use

of video games may lead to the possible development of IGD. To this end, different prevention options for adolescents have been proposed during the pandemic [170]: (i) promotion by schools of avenues for interaction among students when schools are closed due to lockdowns; (ii) monitoring by parents of gaming patterns (e.g. (iii) promotion of home activities, such as table games or physical exercise at home; (iv) promotion of educational video games instead of mainstream video games; and (v) promotion by clinicians of emotional support and coping strategies to better address the pandemic.

5. IGD treatment

In addition to efforts to prevent problems associated with gaming behavior, attempts to treat IGD have been increasing worldwide, and especially in East Asia, despite the fact that there is still no complete consensus on the definition and assessment of this psychiatric entity [171]. Currently, studies of psychotherapy for IGD outnumber those involving pharmacological approaches 3 to 1 [172].

5.1. Pharmacotherapy

Zajac et al. [173], in their systematic review, found that 5 of the 13 studies focused on treatment for IGD evaluated the efficacy of pharmacotherapy. Zajac et al. [174], included 7 pharmacotherapy studies in another systematic review. The most commonly used drugs for the pharmacological approach to IGD were those psychotropic drugs that are also used to treat ADHD (methylphenidate and atomoxetine) and depressive symptomatology (bupropion and escitalopram). Although it has been suggested that these drugs may be effective in reducing the gaming time and IGD symptomatology and severity, even correlating with a reduction in impulsivity levels [175,176], more empirical evidence is needed to draw more solid conclusions.

5.2. Psychotherapy

5.2.1. Cognitive behavioral therapy

A recent systematic review found 4 studies evaluating the effect of cognitive behavioral therapy (CBT) on IGD [173]. It has been suggested that CBT has an impact on the reduction of weekly gaming hours and IGD symptomatology. King et al. [171] also found in their systematic review that CBT was the psychological intervention with the most empirical evidence. However, the authors noted that one of the central limitations of these studies was that they were ambiguous in explaining the content covered in the various treatment sessions. In addition, Zajac et al. [174] reported in their systematic review that, although the treatments employed focused on CBT principles, the specific type of CBT varied, with studies including standard CBT, and others employing craving-focused CBT, gaming-specific CBT, or CBT with mindfulness strategies.

The meta-analysis of Stevens et al. [177] included 12 studies focusing on CBT for patients with IGD, most of them conducted in Asia. The authors found a significant reduction in IGD and depressive symptomatology at post-test. However, it was not possible to determine the efficacy of CBT in reducing the time spent gaming. In addition, Han et al. [178] suggested that CBT may also have an effect in reducing anxious symptomatology comorbid to IGD, as well as impulsivity levels and social avoidance. However, although CBT appears to be a promising intervention for this disorder, more evidence is needed.

In this vein, Young & Brand [179] propose the possible use of CBT for Internet addiction (CBT-IA) for IGD. This intervention, initially developed for Internet addiction, would be used by linking different elements, such as maladaptive cognitions, to online games. The intervention consists of three stages: behavioral modification, cognitive restructuring and harm reduction.

5.2.2. Other approaches

New alternative approaches to standard CBT have emerged in recent years, possibly with the aim of reducing dropout rates, as well as making the interventions more attractive to the target of IGD patients, who are mostly adolescents [180]. According to Wendt et al. [181], these interventions for adolescents with IGD should, in addition to addressing elements of CBT (e.g., reduction of gaming times), generate a structured, adaptive and supportive therapeutic environment.

Four studies were included in the systematic review by Zajac et al. [173] analyzing other approaches to IGD. Family therapy, eclectic psychotherapy (including cognitive behavioral therapy strategies, family therapy, solution-focused therapy and motivational interviewing), the self-discovery camp for adolescents (composed of cognitive behavioral therapy sessions and personal counseling, among others) and a writing and speaking course were tested in these studies. The studies reported a decrease in gaming time and parent-reported IGD symptomatology.

Multidimensional family therapy and family therapy as usual, whose effectiveness has been previously tested in other disorders, appear to be a promising option for treating IGD in adolescents [182,183]. This intervention focuses on the role of the family in both the development and management of adolescent disorders. Kim & Noh [184] highlight that family-based interventions are effective in reducing both online game playing time and the severity of IGD.

Other manualized treatment programs have been proposed to address IGD, such as OMPRIIS and PIPATIC. On the one hand, OMPRIIS combines motivational interviews, elements from CBT, social counseling, media education and Internet-related addiction therapy [185]. On the other hand, PIPATIC is an intervention for adolescents between 12 and 18 years of age consisting of six modules (psychoeducational, treatment as usual, intrapersonal, interpersonal, family intervention, and development of a new lifestyle) aimed at reducing IGD symptomatology [186]. PIPATIC has been shown to be effective for both IGD and comorbid symptomatology, as well as for the improvement of family and interpersonal relationships [187]. Another example is the use of equine-assisted activities and therapies, which have been suggested to have a direct impact on the attachment of adolescents with IGD and insecure attachment issues, which would consequently reduce the severity of IGD symptomatology [188].

The therapeutic residential camp (self-discovery camp) is an intensive treatment for adolescents with IGD developed in South Korea that includes a combination of different therapies, such as CBT, psychoeducation, medical lectures, personal counseling and workshops. This intervention has shown improvements in total gaming time, as well as in problem recognition and self-efficacy towards positive change [189].

Mindfulness has also been proposed as an approach to IGD, showing reductions in maladaptive cognitions associated with gaming behavior which, in turn, appear to have an effect on reducing IGD severity and craving [190,191].

Finally, other approaches have been considered, such as transcranial direct current stimulation [192,193], high-frequency repetitive transcranial magnetic stimulation [194] or virtual reality therapy [180], although their efficacy has not yet been explored in depth.

5.3. Challenges of treatment studies

King et al. [171] highlighted, after their CONSORT evaluation, that there are still unresolved issues and limitations regarding the different treatments proposed for IGD, such as: (1) the optimal duration of the intervention; (2) the differences in efficacy between group and individual format; (3) the lack of control groups; (4) the inconsistencies regarding the assessment of treatment outcome; and (5) the lack of follow-up.

6. Limitations and future studies

The main limitation in the study of IGD is the lack of a consensual theoretical model and, consequently, the lack of diagnostic criteria and cut-off points whose validity has been tested. Therefore, future studies, especially longitudinal studies, as well as international multicenter collaborations, should deepen the clinical validity of the current classification systems and possible consensus cut-off points, as well as the selection of an adequate psychometric instrument that allows the comparison of results between studies. To remedy the lack of literature focused on this disorder, different authors have proposed the following [195–198]:

- 1) To explore the risk factors for the development of IGD in depth (especially genetic, social, and personality factors), and its impact on quality of life and health. Moreover, to identify possible biomarkers in order to improve detection and intervention.
- 2) Although there is some research on how different genres of games (from strategic simulation games to arcade games) have differing levels of addictive potential, it is essential to continue to explore the new video games that are constantly being developed and which ones include structural characteristics with greater addictive potential.
- 3) To carry out more studies focused on the clinical population, instead of the general population.
- 4) To design assessment instruments that are age and culturally appropriate in order to explore the disorder contemplating these essential factors.
- 5) To focus on the comorbidities of IGD in order to discover whether it is a separate entity or a symptom of other psychiatric conditions, such as ADHD, depression or anxiety.
- 6) To develop longitudinal studies, especially randomized controlled trials, in order to deepen the clinical course of the disorder to know, for example, whether it is a continuous or intermittent pathology, or how long it takes to develop, and to study the effectiveness of different therapeutic approaches.

7. Conclusions

Understanding the precise conceptualization of IGD has important implications for its assessment, prevention and treatment. Therefore, these findings support the need to increase the evidence focused on this disorder in order to reach a scientific consensus about the most effective theoretical model, psychometric instrument and treatment plan.

Funding

This research was funded by Ministerio de Ciencia, Innovación y Universidades (grant RTI2018-101837-B-I00). FIS PI20/132, FIS PI17/01167 received aid from the Instituto Salud Carlos III (Ministerio de Sanidad, Servicios Sociales e Igualdad). It was also funded by the Delegación del Gobierno para el Plan Nacional sobre Drogas (2019I47 and 2021I031), CIBERObn-Fisiología Obesidad y Nutrición is an initiative of ISCIII. We thank CERCA Programme/Generalitat de Catalunya for institutional support. Fondo Europeo de Desarrollo Regional (FEDER) “Una manera de hacer Europa”/“A way to build Europe”. Gemma Mestre-Bach was supported by a postdoctoral grant of FUNCIVA.

Conflicts of interest

F.F.-A. received consultancy honoraria from Novo Nordisk and editorial honoraria as EIC from Wiley. The rest of the authors declare no conflict of interest. The funders had no role in the design of the study; the collection, analyses, or interpretation of data; the writing of the manuscript, or the decision to publish the results.

References

- [1] American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders 5 edition (DSM-5). Washington, DC: American Psychiatric Association; 2013.
- [2] World Health Organization. International Statistical Classification of Diseases and Related Health Problems. 11th ed. 2019.
- [3] Kardefelt-Winther D. Conceptualizing Internet use disorders: addiction or coping process? *Psychiatry Clin Neurosci* 2017;71:459–66.
- [4] Starcevic V. Internet Gaming Disorder: inadequate diagnostic criteria wrapped in a constraining conceptual model: commentary on: chaos and confusion in DSM-5 diagnosis of Internet Gaming Disorder: issues, concerns, and recommendations for clarity in the field (Kuss et al.). *J Behav Addict* 2017;6:110–3.
- [5] Markey PM, Ferguson CJ. Internet gaming addiction: disorder or moral panic? *Am J Psychiatry* 2017;174:195–6.
- [6] Dullur P, Hay P. Problem internet use and internet gaming disorder: a survey of health literacy among psychiatrists from Australia and New Zealand. *Australas Psychiatry* 2017;25:140–5.
- [7] King DL, Delfabbro PH, Potenza MN, Demetrovics Z, Billieux J, Brand M. Internet gaming disorder should qualify as a mental disorder. *Aust N Z J Psychiatry* 2018;52:615–7.
- [8] Castro-Calvo J, King DL, Stein DJ, Brand M, Carmi L, Chamberlain SR, et al. Expert appraisal of criteria for assessing gaming disorder: an international Delphi study. *Addiction* 2021;116:2463–75.
- [9] Chen KH, Oliffe JL, Kelly MT. Internet gaming disorder: an emergent health issue for men. *Am J Mens Health* 2018;12:1151–9.
- [10] Paulus FW, Ohmann S, von Gontard A, Popow C. Internet gaming disorder in children and adolescents: a systematic review. *Dev Med Child Neurol* 2018;60:645–59.
- [11] Stevens MWR, Dorstyn D, Delfabbro PH, King DL. Global prevalence of gaming disorder: a systematic review and meta-analysis. *Aust N Z J Psychiatry* 2021;55:553–68.
- [12] Fam JY. Prevalence of internet gaming disorder in adolescents: a meta-analysis across three decades. *Scand J Psychol* 2018;59:524–31.
- [13] Darvesh N, Radhakrishnan A, Lachance CC, Nincic V, Sharpe JP, Ghassemi M, et al. Exploring the prevalence of gaming disorder and Internet gaming disorder: a rapid scoping review. *Syst Rev* 2020;9:1–10.
- [14] Pan YC, Chiu YC, Lin YH. Systematic review and meta-analysis of epidemiology of internet addiction. *Neurosci Biobehav Rev* 2020;118:612–22.
- [15] van Rooij AJ, Van Looy J, Billieux J. Internet Gaming Disorder as a formative construct: implications for conceptualization and measurement. *Psychiatry Clin Neurosci* 2017;71:445–58.
- [16] King DL, Billieux J, Carragher N, Delfabbro PH. Face validity evaluation of screening tools for gaming disorder: scope, language, and overpathologizing issues. *J Behav Addict* 2020;9:1–13.
- [17] Bernaldo-de-Quirós M, Labrador-Méndez M, Sánchez-Iglesias I, Labrador F. Measurement instruments of Internet gaming disorder in adolescents and young people according to DSM-5 criteria: a systematic review. *Adicciones* 2020;32:63–76.
- [18] King DL, Chamberlain SR, Carragher N, Billieux J, Stein D, Mueller K, et al. Screening and assessment tools for gaming disorder: a comprehensive systematic review. *Clin Psychol Rev* 2020;77:101831.
- [19] Yu SM, Pesigan IJA, Zhang MX, Wu AMS. Psychometric validation of the internet gaming disorder-20 test among Chinese middle school and university students. *J Behav Addict* 2019;8:295–305.
- [20] Kim BN. Korean validation of the internet gaming disorder-20 test. *Cyberpsychol Behav Soc Netw* 2019;22:271–6.
- [21] Hawi NS, Samaha M. Validation of the Arabic version of the internet gaming disorder-20 test. *Cyberpsychol Behav Soc Netw* 2017;20:268–72.
- [22] Plessis C, Altintas E, Romo L, Guerrien A. Validation française d’une échelle de mesure de l’utilisation problématique des jeux vidéo en ligne: l’Internet Gaming Disorder-20: French validation of a scale evaluating Internet Gaming Disorder: the Internet Gaming Disorder-20. *Can J Psychiatry* 2021;66:1051–8.
- [23] Männikkö N, Ruotsalainen H, Tolvanen A, Kääriäinen M. Psychometric properties of the Internet Gaming Disorder Test (IGDT-10) and problematic gaming behavior among Finnish vocational school students. *Scand J Psychol* 2019;60:252–60.
- [24] Király O, Bóthe B, Ramos-Díaz J, Rahimi-Movaghar A, Lukavská K, Hrabec O, et al. Ten-Item Internet Gaming Disorder Test (IGDT-10): measurement invariance and cross-cultural validation across seven language-based samples. *Psychol Addict Behav* 2019;33:91–103.
- [25] Király O, Slezcka P, Pontes HM, Urbán R, Griffiths MD, Demetrovics Z. Validation of the Ten-Item Internet Gaming Disorder Test (IGDT-10) and evaluation of the nine DSM-5 Internet Gaming Disorder criteria. *Addict Behav* 2017;64:253–60.
- [26] Chiu YC, Pan YC, Lin YH. Chinese adaptation of the Ten-Item Internet Gaming Disorder Test and prevalence estimate of Internet gaming disorder among adolescents in Taiwan. *J Behav Addict* 2018;7:719–26.
- [27] Wu TY, Lin CY, Årestedt K, Griffiths MD, Broström A, Pakpour AH. Psychometric validation of the Persian nine-item Internet Gaming Disorder Scale – Short Form: does gender and hours spent online Gaming affect the interpretations of item descriptions? *J Behav Addict* 2017;6:256–63.
- [28] Chen IH, Ahorsu DK, Pakpour AH, Griffiths MD, Lin CY, Chen CY. Psychometric properties of three simplified Chinese online-related addictive behavior instruments among mainland Chinese primary school students. *Front Psych* 2020;11:1–8.

- [29] Qin L, Cheng L, Hu M, Liu Q, Tong J, Hao W, et al. Clarification of the cut-off score for nine-item Internet Gaming Disorder Scale–Short Form (IGDS9-SF) in a Chinese context. *Front Psych* 2020;11:1–6.
- [30] Schivinski B, Brzozowska-Woś M, Buchanan EM, Griffiths MD, Pontes HM. Psychometric assessment of the Internet Gaming Disorder diagnostic criteria: an Item Response Theory study. *Addict Behav Rep* 2018;8:176–84.
- [31] Monacis L, De Palo V, Griffiths MD, Sinatra M. Validation of the Internet Gaming Disorder Scale - Short-Form (IGDS9-SF) in an Italian-speaking sample. *J Behav Addict* 2016;5:683–90.
- [32] Beranuy M, Machimbarrena JM, Vega-osés MA, Carbonell X, Griffiths MD, Pontes HM, et al. Spanish validation of the internet gaming disorder scale–short form (IGDS9-SF): prevalence and relationship with online gambling and quality of life. *Int J Environ Res Public Health* 2020;17.
- [33] Maldonado-Murciano L, Pontes HM, Griffiths MD, Barrios M, Gómez-Benito J, Guilera G. The Spanish version of the internet gaming disorder scale-short form (IGDS9-SF): further examination using item response theory. *Int J Environ Res Public Health* 2020;17:1–14.
- [34] Sanchez-Iglesias I, Bernaldo-De-Quiros M, Labrador FJ, Estupina Puig FJ, Labrador M, Fernandez-Arias I. Spanish Validation and Scoring of the Internet Gaming Disorder Scale – Short-Form (IGDS9-SF). *Span J Psychol* 2020;23:1–11.
- [35] Severo RB, Barbosa APPN, Fouchy DRC, Coelho FM da C, Pinheiro RT, de Figueiredo VLM, et al.. Development and psychometric validation of Internet Gaming Disorder Scale-Short-Form (IGDS9-SF) in a Brazilian sample. *Addict Behav* 2020;103:106191.
- [36] Donadon MF, Chagas MHN, Apolinário-Da-silva TD, Okino ETK, Hallak JEC, Nicoletti EA, et al. Cross-cultural adaptation of the internet gaming disorder scale – short form (IGDS9-SF) to the Brazilian context. *Trends Psychiatry Psychother* 2020;42:262–6.
- [37] Ling SL, Jaafar NRN, Tan KA, Bahar N, Baharudin A, Tajjudin AIA. Psychometric properties of the malay version of the internet gaming disorder scale-short form (Igd9-sf-m): evidence from a sample of Malaysian undergraduates. *Int J Environ Res Public Health* 2021;18:1–9.
- [38] Evren C, Dalbudak E, Topcu M, Kutlu N, Evren B, Pontes HM. Psychometric validation of the Turkish nine-item Internet Gaming Disorder Scale–Short Form (IGDS9-SF). *Psychiatry Res* 2018;265:349–54.
- [39] Kim BN, Ko H. Psychometric properties of the Nine-Item Korean internet gaming disorder scale: short form. *Cyberpsychol Behav Soc Netw* 2020;23:854–9.
- [40] Lei W, Liu K, Zeng Z, Liang X, Huang C, Gong K, et al. The psychometric properties of the Chinese version internet gaming disorder scale. *Addict Behav* 2020;106:106392.
- [41] Sigerson L, Li AYL, Cheung MWL, Luk JW, Cheng C. Psychometric properties of the Chinese Internet Gaming Disorder Scale. *Addict Behav* 2017;74:20–6.
- [42] Paschke K, Sack PM, Thomasius R. Validity and psychometric properties of the internet gaming disorder scale in three large independent samples of children and adolescents. *Int J Environ Res Public Health* 2021;18:1–15.
- [43] van Rooij AJ, Schoenmakers TM, van de Mheen D. Clinical validation of the C-VAT 2.0 assessment tool for gaming disorder: a sensitivity analysis of the proposed DSM-5 criteria and the clinical characteristics of young patients with ‘video game addiction’. *Addict Behav* 2017;64:269–74.
- [44] Lin CY, Imani V, Broström A, Årestedt K, Pakpour AH, Griffiths MD. Evaluating the psychometric properties of the 7-item persian game addiction scale for Iranian adolescents. *Front Psychol* 2019;10:1–13.
- [45] Liu Y, Wang Q, Jou M, Wang B, An Y, Li Z. Psychometric properties and measurement invariance of the 7-item game addiction scale (GAS) among Chinese college students. *BMC Psychiatry* 2020;20:1–11.
- [46] Khazaal Y, Breivik K, Billieux J, Zullino D, Thorens G, Achab S, et al. Game addiction scale assessment through a nationally representative sample of young adult men: item response theory graded-response modeling. *J Med Internet Res* 2018;20:e10058.
- [47] Chen JH, Zhang MX, Ko CH, Tong KK, Yu SM, Sou EKL, et al. The development of a screening tool for chinese disordered gamers: the chinese internet gaming disorder checklist (c-igdc). *Int J Environ Res Public Health* 2020;17.
- [48] Lau JTF, Dang L, Cheung RYH, Zhang MX, Chen JH, Wu AMS. The psychometric properties of the Revised Illness Perception Questionnaire (IPQ-R) regarding Internet gaming disorder in a general population of Chinese adults. *J Behav Addict* 2020;9:654–63.
- [49] Higuchi S, Osaki Y, Kinjo A, Mihara S, Maezono M, Kitayuguchi T, et al. Development and validation of a nine-item short screening test for ICD-11 gaming disorder (GAMES test) and estimation of the prevalence in the general young population. *J Behav Addict* 2021;10:263–80.
- [50] Yang X, Jiang X, Wu AMS, Ma L, Cai Y, Wong KM, et al. Validation of the internet gaming disorder symptoms checklist based on the fifth edition of the diagnostic and statistical manual of mental disorders in chinese adolescents. *Child Psychiatry Hum Dev* 2021:1–8.
- [51] Ko CH, Chen SH, Wang CH, Tsai WX, Yen JY. The clinical utility of the chen internet addiction scale—gaming version, for internet gaming disorder in the DSM-5 among young adults. *Int J Environ Res Public Health* 2019;16.
- [52] Basu S, Sharma P, Rustagi R, Sharma R, Sharma N. Measuring addiction to internet gaming among Indian medical students: development and preliminary psychometric properties of a new scale. *Ind Psychiatry J* 2020;29:33–7.
- [53] Irmak AY, Çelikalp Ü, Aydın GÖ, Metinoglu M. Development and validation of the gaming disorder scale–family form. *J Psychosoc Nurs Ment Health Serv* 2021: 1–8.
- [54] Kim BN, Kang HS. Korean validation of the Motives for Online Gaming Questionnaire: focusing on its factor structure and incremental validity. *Addict Behav* 2021;122:107019.
- [55] Wu AMS, Lai MHC, Yu S, Lau JTF, Lei MW. Motives for online gaming questionnaire: its psychometric properties and correlation with Internet gaming disorder symptoms among Chinese people. *J Behav Addict* 2017;6:11–20.
- [56] Yu Y, Mo PKH, Zhang J, Li J, Lau JTF. Validation of the chinese version of the revised internet gaming cognition scale among adolescents in China: maladaptive cognitions as potential determinants of internet gaming disorder. *Int J Environ Res Public Health* 2020;17.
- [57] Hawi NS, Samaha M, Griffiths MD. The digital addiction scale for children: development and validation. *Cyberpsychol Behav Soc Netw* 2019;22:771–8.
- [58] Sarda E, Begue L, Bry C, Gentile D. Internet gaming disorder and well-being: a scale validation. *Cyberpsychol Behav Soc Netw* 2016;19:674–9.
- [59] Koo HJ, Han DH, Park SY, Kwon JH. The structured clinical interview for DSM-5 internet gaming disorder: development and validation for diagnosing IGD in adolescents. *Psychiatry Investig* 2017;14:21–9.
- [60] Sprong ME, Griffiths MD, Lloyd DP, Paul E, Buono FD. Comparison of the video game functional assessment-revised (VGFA-R) and internet gaming disorder test (IGD-20). *Front Psychol* 2019;10:1–7.
- [61] Stavropoulos V, Beard C, Griffiths MD, Buleigh T, Gomez R, Pontes HM. Measurement invariance of the internet gaming disorder scale–short-form (IGDS9-SF) between Australia, the USA, and the UK. *Int J Ment Health Addict* 2018;16:377–92.
- [62] Pontes HM, Stavropoulos V, Griffiths MD. Measurement invariance of the internet gaming disorder scale–short-form (IGDS9-SF) between the United States of America, India and the United Kingdom. *Psychiatry Res* 2017;257:472–8.
- [63] Yilmaz E, Griffiths MD, Kan A. Development and validation of videogame addiction scale for children (VASC). *Int J Ment Health Addict* 2017;15:869–82.
- [64] Jo SJ, Yim HW, Lee HK, Lee HC, Choi JS, Baek KY. The internet game use-elicited symptom screen proved to be a valid tool for adolescents aged 10–19 years. *Acta Paediatr Int J Paediatr* 2018;107:511–6.
- [65] Ryu H, Lee JY, Choi AR, Chung SJ, Park M, Bhang SY, et al. Application of diagnostic interview for internet addiction (DIA) in clinical practice for Korean adolescents. *J Clin Med* 2019;8.
- [66] Besser B, Rumpf HJ, Bischof A, Meerkerk GJ, Higuchi S, Bischof G. Internet-related disorders: development of the short compulsive internet use scale. *Cyberpsychol Behav Soc Netw* 2017;20:709–17.
- [67] Gmel G, Khazaal Y, Studer J, Baggio S, Marmet S. Development of a short form of the compulsive internet use scale in Switzerland. *Int J Methods Psychiatr Res* 2019;28:1–11.
- [68] Schluter MG, Hodgins DC, Wolfe J, Wild TC. Can one simple questionnaire assess substance-related and behavioural addiction problems? Results of a proposed new screener for community epidemiology. *Addiction* 2018;113:1528–37.
- [69] Leung H, Pakpour AH, Strong C, Lin YC, Tsai MC, Griffiths MD, et al. Measurement invariance across young adults from Hong Kong and Taiwan among three internet-related addiction scales: Bergen Social Media Addiction Scale (BSMAS), Smartphone Application-Based Addiction Scale (SABAS), and Internet Gaming Disorder Scale-Short. *Addict Behav* 2020;101:105969.
- [70] Chen IH, Strong C, Lin YC, Tsai MC, Leung H, Lin CY, et al. Time invariance of three ultra-brief internet-related instruments: smartphone Application-Based Addiction Scale (SABAS), Bergen Social Media Addiction Scale (BSMAS), and the nine-item Internet Gaming Disorder Scale- Short Form (IGDS-SF9) (Study Part B). *Addict Behav* 2020;101:105960.
- [71] Wang HY, Cheng C. Psychometric evaluation and comparison of two gaming disorder measures derived from the DSM-5 and ICD-11 frameworks. *Front Psych* 2020;11:1–11.
- [72] Kuss DJ, Griffiths MD, Pontes HM. DSM-5 diagnosis of Internet Gaming disorder: some ways forward in overcoming issues and concerns in the Gaming studies field: response to the commentaries. *J Behav Addict* 2017;6:133–41.
- [73] Brand M, Young KS, Laier C. Prefrontal control and Internet addiction: a theoretical model and review of neuropsychological and neuroimaging findings. *Front Hum Neurosci* 2014;8:1–13.
- [74] Dong G, Potenza MN. A cognitive-behavioral model of Internet gaming disorder: theoretical underpinnings and clinical implications. *J Psychiatr Res* 2014;58: 7–11.
- [75] Brand M, Young KS, Laier C, Wölfling K, Potenza MN. Integrating psychological and neurobiological considerations regarding the development and maintenance of specific Internet-use disorders: an Interaction of Person-Affect-Cognition-Execution (I-PACE) model. *Neurosci Biobehav Rev* 2016;71:252–66.
- [76] Yen JY, Lin PC, Lin HC, Lin PY, Chou WP, Ko CH. Association of Internet gaming disorder with catechol-O-methyltransferase: role of impulsivity and fun-seeking. *Kaohsiung J Med Sci* 2021:1–7.
- [77] Katteine E, Schmidt H, Brandt D, Bischof G, Bischof A, Besser B, et al. Association of increased impulsiveness and internet use disorder in adolescents and young adults with different main activities on the internet. *Z Kinder Jugendpsychiatr Psychother* 2021;0:1–8.
- [78] Chun JW, Park CH, Kim JY, Choi J, Cho H, Jung DJ, et al. Altered core networks of brain connectivity and personality traits in internet gaming disorder. *J Behav Addict* 2020;9:298–311.
- [79] Efrati Y, Kolubinski DC, Marino C, Spada MM. Modelling the contribution of metacognitions, impulsiveness and thought suppression to behavioural addictions in adolescents. *Int J Environ Res Public Health* 2021;18.
- [80] Ryu H, Lee JY, Choi A, Park S, Kim DJ, Choi JS. The relationship between impulsivity and internet gaming disorder in young adults: mediating effects of interpersonal relationships and depression. *Int J Environ Res Public Health* 2018; 15:1–11.

- [81] Hu J, Zhen S, Yu C, Zhang Q, Zhang W. Sensation seeking and online gaming addiction in adolescents: a moderated mediation model of positive affective Associations and Impulsivity. *Front Psychol* 2017;8:1–8.
- [82] Tian Y, Yu C, Lin S, Lu J, Liu Y, Zhang W. Sensation seeking, deviant peer affiliation, and internet gaming addiction among Chinese adolescents: the moderating effect of parental knowledge. *Front Psychol* 2019;9:1–7.
- [83] Shin D, Choi AR, Lee J, Chung SJ, Kim B, Park M, et al. The mediating effects of affect on associations between impulsivity or resilience and internet gaming disorder. *J Clin Med* 2019;8:1–11.
- [84] Wölfling K, Duvén E, Wejbera M, Beutel ME, Müller KW. Discounting delayed monetary rewards and decision making in behavioral addictions – a comparison between patients with gambling disorder and internet gaming disorder. *Addict Behav* 2020;108.
- [85] Raiha S, Yang G, Wang L, Dai W, Wu H, Meng G, et al. Altered reward processing system in internet gaming disorder. *Front Psych* 2020;11:1–13.
- [86] Xiang H, Tian X, Zhou Y, Chen J, Potenza MN, Zhang Q. The relationship between behavioral inhibition and behavioral activation systems, impulsiveness, and internet gaming disorder among students of different ages. *Front Psych* 2021;11:1–6.
- [87] Müller KW, Werthmann J, Beutel ME, Wölfling K, Egloff B. Maladaptive personality traits and their interaction with outcome expectancies in gaming disorder and internet-related disorders. *Int J Environ Res Public Health* 2021;18.
- [88] Liao Z, Huang Q, Huang S, Tan L, Shao T, Fang T, et al. Prevalence of internet gaming disorder and its association with personality traits and gaming characteristics among Chinese adolescent gamers. *Front Psych* 2020;11:1–8.
- [89] Lee YS, Son JH, Park JH, Kim SM, Kee BS, Han DH. The comparison of temperament and character between patients with internet gaming disorder and those with alcohol dependence. *J Ment Health* 2017;26:242–7.
- [90] Granero R, Fernández-aranda F, Demetrovics Z, Ayala-rojas RE, Gómez-peña M, Moragas L, et al. Profile of treatment – seeking gaming disorder patients: a network perspective. *J Gambli Stud* 2021.
- [91] González-Bueso V, Santamaría JJ, Oliveras I, Fernández D, Montero E, Baño M, et al. Internet gaming disorder clustering based on personality traits in adolescents, and its relation with comorbid psychological symptoms. *Int J Environ Res Public Health* 2020;17.
- [92] Mallorquí-Bagué N, Fernández-Aranda F, Lozano-Madrid M, Granero R, Mestre-Bach G, Baño M, et al. Internet gaming disorder and online gambling disorder: clinical and personality correlates. *J Behav Addict* 2017;6:669–77.
- [93] Chung W, Sun CK, Tsai IT, Hung KC, Chiu HJ, Tzang RF, et al. A systematic review and meta-analysis on the clinical implications of probability discounting among individuals with Internet gaming disorder. *Sci Rep* 2021;11:1–10.
- [94] Wang Y, Wu L, Wang L, Zhang Y, Du X, Dong G. Impaired decision-making and impulse control in Internet gaming addicts: evidence from the comparison with recreational Internet game users. *Addict Biol* 2017;22:1610–21.
- [95] Liu L, Xue G, Potenza MN, Zhang JT, Yao YW, Xia CC, et al. Dissociable neural processes during risky decision-making in individuals with Internet-gaming disorder. *NeuroImage Clin* 2017;14:741–9.
- [96] Sen Yan W, Chen RT, Liu MM, Zheng DH. Monetary reward discounting, inhibitory control, and trait impulsivity in young adults with internet gaming disorder and nicotine dependence. *Front Psych* 2021;12:1–14.
- [97] Chen J, Li X, Zhang Q, Zhou Y, Wang R, Tian C, et al. Impulsivity and response inhibition related brain networks in adolescents with internet gaming disorder: a preliminary study utilizing resting-state fMRI. *Front Psych* 2021;11:1–11.
- [98] Yao YW, Zhang JT, Fang XY, Liu L, Potenza MN. Reward-related decision-making deficits in internet gaming disorder: a systematic review and meta-analysis. *Addiction* 2021;117:19–32.
- [99] Kräplin A, Scherbaum S, Kraft EM, Rehbein F, Bühringer G, Goshcke T, et al. The role of inhibitory control and decisionmaking in the course of Internet gaming disorder. *J Behav Addict* 2021;9:990–1001.
- [100] Yen JY, Yeh YC, Wang PW, Liu TL, Chen YY, Ko CH. Emotional regulation in young adults with internet gaming disorder. *Int J Environ Res Public Health* 2018; 15:1–11.
- [101] Lin PY, Lin HC, Lin PC, Yen JY, Ko CH. The association between emotional regulation and internet gaming disorder. *Psychiatry Res* 2020;289:113060.
- [102] Wu L, Potenza MN, Zhou N, Kober H, Shi X, Yip SW, et al. A role for the right dorsolateral prefrontal cortex in enhancing regulation of both craving and negative emotions in internet gaming disorder: a randomized trial. *Eur Neuropsychopharmacol* 2021;36:29–37.
- [103] Liu S, Yu C, Conner BT, Wang S, Lai W, Zhang W. Autistic traits and internet gaming addiction in Chinese children: the mediating effect of emotion regulation and school connectedness. *Res Dev Disabil* 2017;68:122–30.
- [104] Che D, Hu J, Zhen S, Yu C, Li B, Chang X, et al. Dimensions of emotional intelligence and online gaming addiction in adolescence: the indirect effects of two facets of perceived stress. *Front Psychol* 2017;8:1–8.
- [105] Dang DL, Zhang MX, Leong KKH, Anise MSW. The predictive value of emotional intelligence for internet gaming disorder: a 1-year longitudinal study. *Int J Environ Res Public Health* 2019;16:12–5.
- [106] Ko CH, Liu TL, Wu HC, Yeh YC, Tsai WW, Yen JY. Psychiatric comorbidities and emotional intelligence in internet gaming disorder: attention deficit hyperactivity disorder, major depressive disorder, generalized anxiety disorder, and social anxiety disorder. *Psychiatry Clin Neurosci* 2021:1–2.
- [107] Wichstrøm L, Stenseng F, Belsky J, von Soest T, Hygen BW. Symptoms of internet gaming disorder in youth: predictors and comorbidity. *J Abnorm Child Psychol* 2019;47:71–83.
- [108] González-Bueso V, Santamaría JJ, Fernández D, Merino L, Montero E, Ribas J. Association between internet gaming disorder or pathological video-game use and comorbid psychopathology: a comprehensive review. *Int J Environ Res Public Health* 2018;15:1–20.
- [109] González-Bueso V, Santamaría JJ, Fernández D, Merino L, Montero E, Jiménez-Murcia S, et al. Internet gaming disorder in adolescents: personality, psychopathology and evaluation of a psychological intervention combined with parent psychoeducation. *Front Psychol* 2018;9.
- [110] Ohayon MM, Roberts L. Internet gaming disorder and comorbidities among campus-dwelling U.S. university students. *Psychiatry Res* 2021;302:114043.
- [111] Hygen BW, Skalická V, Stenseng F, Belsky J, Steinsbekk S, Wichstrøm L. The co-occurrence between symptoms of internet gaming disorder and psychiatric disorders in childhood and adolescence: prospective relations or common causes? *J Child Psychol Psychiatry Allied Discip* 2020;61:890–8.
- [112] De Pasquale C, Sciacca F, Martinielli V, Chiappelli M, Dinaro C, Hichy Z. Relationship of internet gaming disorder with psychopathology and social adaptation in Italian young adults. *Int J Environ Res Public Health* 2020;17:1–8.
- [113] Ha Park J, Sun Hong J, Hyun Han D, Joon Min K, Sik Lee Y, Seok Kee B, et al. Comparison of QEEG findings between adolescents with attention deficit hyperactivity disorder (ADHD) without comorbidity and ADHD comorbid with internet gaming disorder. *J Korean Med Sci* 2017;32:514–21.
- [114] Yen JY, Liu TL, Wang PW, Chen CS, Yen CF, Ko CH. Association between Internet gaming disorder and adult attention deficit and hyperactivity disorder and their correlates: impulsivity and hostility. *Addict Behav* 2017;64:308–13.
- [115] Chen C, Dai S, Shi L, Shen Y, Ou J. Associations between attention deficit/hyperactivity disorder and internet gaming disorder symptoms mediated by depressive symptoms and hopelessness among college students. *Neuropsychiatr Dis Treat* 2021;17:2775–82.
- [116] Evren C, Evren B, Dalbudak E, Topcu M, Kutlu N. Relationships of Internet addiction and Internet gaming disorder symptom severities with probable attention deficit/hyperactivity disorder, aggression and negative affect among university students. *ADHD Atten Deficit Hyperact Disord* 2019;11:413–21.
- [117] Kim M, Kim D, Bae S, Han DH, Jeong B. Aberrant structural network of comorbid attention deficit/hyperactivity disorder is associated with addiction severity in internet gaming disorder. *NeuroImage Clin* 2020;27:102263.
- [118] Dullur P, Krishnan V, Diaz AM. A systematic review on the intersection of attention-deficit hyperactivity disorder and gaming disorder. *J Psychiatr Res* 2021;133:212–22.
- [119] Lee J, Bae S, Kim BN, Han DH. Impact of attention-deficit/hyperactivity disorder comorbidity on longitudinal course in Internet gaming disorder: a 3-year clinical cohort study. *J Child Psychol Psychiatry Allied Discip* 2021;62:1110–9.
- [120] Stavropoulos V, Adams BLM, Beard CL, Dumble E, Trawley S, Gomez R, et al. Associations between attention deficit hyperactivity and internet gaming disorder symptoms: is there consistency across types of symptoms, gender and countries? *Addict Behav Rep* 2019;9:100158.
- [121] Concerto C, Rodolico A, Avanzato C, Fusar-Poli L, Signorelli MS, Battaglia F, et al. Autistic traits and attention-deficit hyperactivity disorder symptoms predict the severity of internet gaming disorder in an Italian adult population. *Brain Sci* 2021; 11.
- [122] Wang R, Yang S, Yan Y, Tian Y, Wang P. Internet gaming disorder in early adolescents: gender and depression differences in a latent growth model. *Healthc* 2021;9.
- [123] Wang HR, Cho H, Kim DJ. Prevalence and correlates of comorbid depression in a nonclinical online sample with DSM-5 internet gaming disorder. *J Affect Disord* 2018;226:1–5.
- [124] Yen JY, Lin HC, Chou WP, Liu TL, Ko CH. Associations among resilience, stress, depression, and internet gaming disorder in young adults. *Int J Environ Res Public Health* 2019;16.
- [125] Bonnaire C, Baptista D. Internet gaming disorder in male and female young adults: the role of alexithymia, depression, anxiety and gaming type. *Psychiatry Res* 2019;272:521–30.
- [126] Liu L, Yao YW, Li C, Shan R, Zhang JT, Xia CC, et al. The comorbidity between internet gaming disorder and depression: interrelationship and neural mechanisms. *Front Psych* 2018;9:1–10.
- [127] Ostinelli EG, Zangani C, Giordano B, Maestri D, Gambini O, D'Agostino A, et al. Depressive symptoms and depression in individuals with internet gaming disorder: a systematic review and meta-analysis. *J Affect Disord* 2021;284: 136–42.
- [128] Yuan G, Elhai JD, Hall BJ. The influence of depressive symptoms and fear of missing out on severity of problematic smartphone use and Internet gaming disorder among Chinese young adults: a three-wave mediation model. *Addict Behav* 2021;112:106648.
- [129] Lin HC, Yen JY, Lin PC, Ko CH. The frustration intolerance of internet gaming disorder and its association with severity and depression. *Kaohsiung J Med Sci* 2021;37:903–9.
- [130] Stavropoulos V, Vassallo J, Burleigh TL, Gomez R, Colder Carras M. The role of internet gaming in the association between anxiety and depression: a preliminary cross-sectional study. *Asia Pac Psychiatry* 2021;14(2):e12474.
- [131] Yu Y, Yang X, Wang S, Wang H, Chang R, Tsamlag L, et al. Serial multiple mediation of the association between internet gaming disorder and suicidal ideation by insomnia and depression in adolescents in Shanghai, China. *BMC Psychiatry* 2020;20:1–9.
- [132] Na E, Lee H, Choi I, Kim DJ. Comorbidity of internet gaming disorder and alcohol use disorder: a focus on clinical characteristics and gaming patterns. *Am J Addict* 2017;26:326–34.
- [133] Wartberg L, Kammerl R. Empirical relationships between problematic alcohol use and a problematic use of video games, social media and the internet and their

- associations to mental health in adolescence. *Int J Environ Res Public Health* 2020;17:1–11.
- [134] Horváth Z, Király O, Demetrovics Z, Németh Á, Várnai D, Urbán R. Polysubstance use is positively associated with gaming disorder symptom severity: a latent class analytical study. *Eur Addict Res* 2021;28:12–22.
- [135] Wong HY, Mo HY, Potenza MN, Chan MNM, Lau WM, Chui TK, et al. Relationships between severity of internet gaming disorder, severity of problematic social media use, sleep quality and psychological distress. *Int J Environ Res Public Health* 2020;17:1–13.
- [136] Pontes HM. Investigating the differential effects of social networking site addiction and Internet gaming disorder on psychological health. *J Behav Addict* 2017;6:601–10.
- [137] Lin CY, Potenza MN, Broström A, Pakpour AH. Internet gaming disorder, psychological distress, and insomnia in adolescent students and their siblings: an actor-partner interdependence model approach. *Addict Behav Rep* 2021;13.
- [138] Kristensen JH, Pallesen S, King DL, Hysing M, Erevik EK. Problematic gaming and sleep: a systematic review and meta-analysis. *Front Psych* 2021;12.
- [139] Murray A, Mannion A, Chen JL, Leader G. Gaming disorder in adults with autism spectrum disorder. *J Autism Dev Disord* 2021;1–8.
- [140] Wang CY, Wu YC, Su CH, Lin PC, Ko CH, Yen JY. Association between Internet gaming disorder and generalized anxiety disorder. *J Behav Addict* 2017;6:564–71.
- [141] Hakami AY, Ahmad RG, Alsharif A, Ashqar A, AlHarbi FA, Sayes M, et al. Prevalence of behavioral addictions and their relationship with stress and anxiety among medical students in Saudi Arabia: a cross-sectional study. *Front Psych* 2021;12:1–9.
- [142] Marino C, Canale N, Vieno A, Caselli G, Scacchi L, Spada MM. Social anxiety and Internet gaming disorder: the role of motives and metacognitions. *J Behav Addict* 2020;9:617–28.
- [143] Andreetta J, Teh MSc J, Burleigh TL, Gomez R, Stavropoulos V. Associations between comorbid stress and Internet Gaming Disorder symptoms: are there cultural and gender variations? *Asia Pac Psychiatry* 2020;12:1–8.
- [144] Rajab AM, Zaghloul MS, Enabi S, Rajab TM, Al-Khani AM, Basalah A, et al. Gaming addiction and perceived stress among Saudi adolescents. *Addict Behav Rep* 2020;11.
- [145] Yuan GF, Shi W, Elhai JD, Montag C, Chang K, Jackson T, et al. Gaming to cope: applying network analysis to understand the relationship between posttraumatic stress symptoms and internet gaming disorder symptoms among disaster-exposed Chinese young adults. *Addict Behav* 2022;124:107096.
- [146] Kardefelt-Winther D. A conceptual and methodological critique of internet addiction research: towards a model of compensatory internet use. *Comput Hum Behav* 2014;31:351–4.
- [147] Aarseth E, Bean AM, Boonen H, Carras MC, Coulson M, Das D, et al. Scholars' open debate paper on the world health organization ICD-11 gaming disorder proposal. *J Behav Addict* 2017;6:267–70.
- [148] Sugaya N, Shirasaka T, Takahashi K, Kanda H. Bio-psychosocial factors of children and adolescents with internet gaming disorder: a systematic review. *Biopsychosoc Med* 2019;13:1–16.
- [149] Schneider LA, King DL, Delfabbro PH. Family factors in adolescent problematic Internet gaming: a systematic review. *J Behav Addict* 2017;6:321–33.
- [150] Wartberg L, Kriston L, Kramer M, Schwedler A, Lincoln T, Kammerl R. Internet gaming disorder in early adolescence: associations with parental and adolescent mental health. *Eur Psychiatry* 2017;43:14–8.
- [151] Bonnaire C, Phan O. Relationships between parental attitudes, family functioning and Internet gaming disorder in adolescents attending school. *Psychiatry Res* 2017;255:104–10.
- [152] Hwang H, Hong J, Kim SM, Han DH. The correlation between family relationships and brain activity within the reward circuit in adolescents with Internet gaming disorder. *Sci Rep* 2020;10:1–9.
- [153] Chen IH, Lee ZH, Dong XY, Gamble JH, Feng HW. The influence of parenting style and time management tendency on internet gaming disorder among adolescents. *Int J Environ Res Public Health* 2020;17:1–16.
- [154] Koning IM, Peeters M, Finkenauer C, Van Den Eijnden RJJM. Bidirectional effects of Internet-specific parenting practices and compulsive social media and Internet game use. *J Behav Addict* 2018;7:624–32.
- [155] Schettler L, Thomasius R, Paschke K. Neural correlates of problematic gaming in adolescents: a systematic review of structural and functional magnetic resonance imaging studies. *Addict Biol* 2021;1–13.
- [156] Kuss DJ, Pontes HM, Griffiths MD. Neurobiological correlates in internet gaming disorder: a systematic literature review. *Front Psych* 2018;9:1–12.
- [157] Choi E, Shin S, Ryu J, Jung K, Hyun Y, Kim JPM. Association of extensive video gaming and cognitive function changes in brain-imaging studies of pro gamers and individuals with gaming disorder: systematic literature review. *JMIR Serious Games* 2021;9:e25793.
- [158] Wang M, Dong H, Zheng H, Du X, Dong GH. Inhibitory neuromodulation of the putamen to the prefrontal cortex in Internet gaming disorder: how addiction impairs executive control. *J Behav Addict* 2020;9:312–24.
- [159] Weinstein A, Lejoyeux M. Neurobiological mechanisms underlying internet gaming disorder. *Dialogues Clin Neurosci* 2020;22:113–26.
- [160] Zhou X, Wu R, Liu C, Kou J, Chen Y, Pontes HM, et al. Higher levels of (Internet) Gaming Disorder symptoms according to the WHO and APA frameworks associate with lower striatal volume. *J Behav Addict* 2020;9:598–605.
- [161] Yao YW, Liu L, Ma SS, Shi XH, Zhou N, Zhang JT, et al. Functional and structural neural alterations in Internet gaming disorder: a systematic review and meta-analysis. *Neurosci Biobehav Rev* 2017;83:313–24.
- [162] King DL, Delfabbro PH. The concept of "harm" in Internet gaming disorder. *J Behav Addict* 2018;1–3.
- [163] Király O, Griffiths MD, King DL, Lee HK, Lee SY, Bányai F, et al. Policy responses to problematic video game use: a systematic review of current measures and future possibilities. *J Behav Addict* 2018;7:503–17.
- [164] King DL, Delfabbro PH, Doh YY, Wu AMS, Kuss DJ, Pallesen S, et al. Policy and prevention approaches for disordered and hazardous gaming and internet use: an international perspective. *Prev Sci* 2018;19:233–49.
- [165] Gentile DA. Thinking more broadly about policy responses to problematic video game use: a response to Király et al. (2018) Commentary on: policy responses to problematic video game use: a systematic review of current measures and future possibilities (Király et al.). *J Behav Addict* 2018;7:536–9.
- [166] Petry NM, Zajac K, Ginley M, Lemmens J, Rumpf HJ, Ko CH, et al. Policy and prevention efforts for gaming should consider a broad perspective Commentary on: policy responses to problematic video game use: a systematic review of current measures and future possibilities (Király et al., 2018). *J Behav Addict* 2018;7:543–7.
- [167] Wegmann E, Brand M. The imperative of integrating empirical and theoretical considerations when developing policy responses to Internet-gaming disorder commentary on: policy responses to problematic video game use: a systematic review of current measures and future possibilities. *J Behav Addict* 2018;7:531–5.
- [168] Kuss DJ. Policy, prevention, and regulation for Internet Gaming Disorder Commentary on: policy responses to problematic video game use: a systematic review of current measures and future possibilities (Király et al., 2018). *J Behav Addict* 2018;7:553–5.
- [169] King DL, Delfabbro PH, Billieux J, Potenza MN. Problematic online gaming and the COVID-19 pandemic. *J Behav Addict* 2020;9:184–6.
- [170] Ko CH, Yen JY. Impact of COVID-19 on gaming disorder: monitoring and prevention. *J Behav Addict* 2020;9:187–9.
- [171] King DL, Delfabbro PH, Wu AMS, Doh YY, Kuss DJ, Pallesen S, et al. Treatment of Internet gaming disorder: an international systematic review and CONSORT evaluation. *Clin Psychol Psychother* 2017;54:123–33.
- [172] King DL, Wölfling K, Potenza MN. Taking gaming disorder treatment to the next level. *JAMA Psychiat* 2020;77:869–70.
- [173] Zajac K, Ginley MK, Chang R, Petry NM. Treatments for internet gaming disorder and internet addiction: a systematic review. *Psychol Addict Behav* 2017;31:979–94.
- [174] Zajac K, Ginley MK, Chang R. Treatments of internet gaming disorder: a systematic review of the evidence. *Expert Rev Neurother* 2020;20:85–93.
- [175] Park JH, Lee YS, Sohn JH, Han DH. Effectiveness of atomoxetine and methylphenidate for problematic online gaming in adolescents with attention deficit hyperactivity disorder. *Hum Psychopharmacol* 2016;31:427–32.
- [176] Bae S, Hong JS, Kim SM, Han DH. Bupropion shows different effects on brain functional connectivity in patients with Internet-based gambling disorder and internet gaming disorder. *Front Psych* 2018;9.
- [177] Stevens MWR, King DL, Dorstyn D, Delfabbro PH. Cognitive-behavioral therapy for Internet gaming disorder: a systematic review and meta-analysis. *Clin Psychol Psychother* 2019;26:191–203.
- [178] Han J, Seo Y, Hwang H, Kim SM, Han DH. Efficacy of cognitive behavioural therapy for internet gaming disorder. *Clin Psychol Psychother* 2020;27:203–13.
- [179] Young KS, Brand M. Merging theoretical models and therapy approaches in the context of internet gaming disorder: a personal perspective. *Front Psychol* 2017;8.
- [180] Zhang M, R. HO. Smartphone applications for immersive and internet gaming disorder. *Technol Health Care* 2017;25:367–72.
- [181] Wendt LM, Austermann MI, Rumpf HJ, Thomasius R, Paschke K. Requirements of a group intervention for adolescents with internet gaming disorder in a clinical setting: a qualitative interview study. *Int J Environ Res Public Health* 2021;18.
- [182] Bonnaire C, Liddle HA, Har A, Nielsen P, Phan O. Why and how to include parents in the treatment of adolescents presenting Internet gaming disorder? *J Behav Addict* 2019;8:201–12.
- [183] Nielsen P, Christensen M, Henderson C, Liddle HA, Croquette-Krokar M, Favez N, et al. Multidimensional family therapy reduces problematic gaming in adolescents: a randomised controlled trial. *J Behav Addict* 2021;10:234–43.
- [184] Kim S, Noh D. The current status of psychological intervention research for internet addiction and internet gaming disorder. *Issues Ment Health Nurs* 2019;40:335–41.
- [185] Dieris-Hirce J, Böttel L, Pape M, Te Wildt BT, Wölfling K, Henningsen P, et al. Effects of an online-based motivational intervention to reduce problematic internet use and promote treatment motivation in internet gaming disorder and internet use disorder (OMPRIS): study protocol for a randomised controlled trial. *BMJ Open* 2021;11:1–12.
- [186] Torres-Rodríguez A, Griffiths MD, Carbonell X. The treatment of internet gaming disorder: a brief overview of the PIPATIC program. *Int J Ment Health Addict* 2018;16:1000–15.
- [187] Torres-Rodríguez A, Griffiths MD, Carbonell X, Oberst U. Treatment efficacy of a specialized psychotherapy program for Internet Gaming Disorder. *J Behav Addict* 2018;7:939–52.
- [188] Kang KD, Jung TW, Park IH, Han DH. Effects of equine-assisted activities and therapies on the affective network of adolescents with internet gaming disorder. *J Altern Complement Med* 2018;24:841–9.
- [189] Sakuma H, Mihara S, Nakayama H, Miura K, Kitayuguchi T, Maezono M, et al. Treatment with the Self-Discovery Camp (SDiC) improves Internet gaming disorder. *Addict Behav* 2017;64:357–62.
- [190] Li W, Garland EL, Howard MO. Therapeutic mechanisms of Mindfulness-Oriented Recovery Enhancement for internet gaming disorder: reducing craving and addictive behavior by targeting cognitive processes. *J Addict Dis* 2018;37:5–13.

- [191] Li W, Garland EL, McGovern P, Brien JEO, Tronnier C, Matthew O. Mindfulness-oriented recovery enhancement for internet gaming disorder in US adults: a stage i randomized controlled trial. *Psychol Addict Behav* 2017;31:393.
- [192] Wu LL, Potenza MN, Zhou N, Kober H, Shi XH, Yip SW, et al. Efficacy of single-session transcranial direct current stimulation on addiction-related inhibitory control and craving: a randomized trial in males with internet gaming disorder. *J Psychiatry Neurosci* 2021;46:E111–8.
- [193] Lee J-Y, Jang JH, Choi AR, Chung SJ, Kim B, Park M, et al. Neuromodulatory effect of transcranial direct current stimulation on resting-state EEG activity in internet gaming disorder: a randomized, double-blind, sham-controlled parallel group trial. *Cereb Cortex Commun* 2021;2:1–10.
- [194] Cuppone D, Perez LJG, Cardullo S, Cellini N, Sarlo M, Soldatesca S, et al. The role of repetitive transcranial magnetic stimulation (rTMS) in the treatment of behavioral addictions: two case reports and review of the literature. *J Behav Addict* 2021;10:361–70.
- [195] Feng W, Ramo DE, Chan SR, Bourgeois JA. Internet gaming disorder: trends in prevalence 1998–2016. *Addict Behav* 2017;75:17–24.
- [196] Kuss DJ, Griffiths MD, Pontes HM. Chaos and confusion in DSM-5 diagnosis of internet gaming disorder: issues, concerns, and recommendations for clarity in the field. *J Behav Addict* 2017;6:103–9.
- [197] Gentile D, Bailey K, Bavelier D, Funk Brockmyer J, Cash H, Coyne S, et al. Internet gaming disorder in children and adolescents. *Pediatrics* 2017;140:S81–5.
- [198] Fineberg NA, Demetrovics Z, Stein DJ, Ioannidis K, Potenza MN, Grünblatt E, et al. Manifesto for a European research network into Problematic Usage of the Internet. *Eur Neuropsychopharmacol* 2018;28:1232–46.